

Supplemental Material to:

**Natalia A Kolganova, Anna M Varizhuk, Roman A Novikov,
Vladimir L Florentiev, Galina E Pozmogova,
Olga F Borisova, Anna K Shchyolkina, Igor P Smirnov,
Dmitry N Kaluzhny, and Edward N Timofeev**

**Anomeric DNA quadruplexes:
Modified thrombin aptamers**

Artificial DNA: PNA & XNA 2013; 5(1)
<http://dx.doi.org/10.4161/adna.28422>

[http://www.landesbioscience.com/journals/artificialdna/
article/28422/](http://www.landesbioscience.com/journals/artificialdna/article/28422/)

Anomeric DNA quadruplexes: modified thrombin aptamers

Natalia A. Kolganova[†], Anna M. Varizhuk^{†‡}, Roman A. Novikov^{†§}, Vladimir L. Florentiev^{1†}, Galina E. Pozmogova[‡], Olga F. Borisova[†], Anna K. Shchyolkina[†], Igor P. Smirnov[‡], Dmitry N. Kaluzhny[†], and Edward N. Timofeev[†]

[†]Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, 119991 Moscow, Russia.

[‡]Institute for Physical-Chemical Medicine of Public Health Ministry, 117312 Moscow, Russia.

[§]Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, 119991 Moscow, Russia.

Corresponding author: edward@eimb.ru

SUPPORTING INFORMATION

¹ Deceased on October 24, 2012.

SUPPLEMENTARY FIGURE LEGENDS

Fig. S1. UV melting profiles of aptamer A12 at 295 nm. (A) Effect of sample annealing rate and KCl concentration on the high-temperature transition. (B) Effect of aptamer concentration on the high-temperature transition.

Fig. S2. Dependence of the T_m value on aptamer concentration in 10 mM sodium cacodylate (pH 7.2) and 100 mM KCl for oligonucleotides A2 and A12. Melting temperatures were determined for slow annealed samples as the minimum of the first derivative of the denaturation profile. Only the major low-temperature transition was analyzed for aptamer A12.

Fig. S3. Thermal difference spectra (80°C vs. 20°C) of TBA and selected modified aptamers in 10 mM sodium cacodylate (pH 7.2) and 100 mM KCl.

Fig. S4. CD spectra of TBA and selected modified aptamers at 20°C in 10 mM sodium cacodylate (pH 7.2) and 100 mM KCl.

Fig. S5. PEG calibration graph and experimental points from diffusion ordered NMR spectroscopy (DOSY) experiments.

Fig. S6. The plot of thrombin activity *vs.* aptamer concentration.

Fig. S7. Calibration curve for converting thrombin time values into thrombin activity. A series of thrombin dilutions was prepared and mixed with blood plasma. Clotting times were measured. Each thrombin time value is an average of three measurements.





FIGURE S1




FIGURE S2




FIGURE S3




FIGURE S4




FIGURE S5




FIGURE S6




FIGURE S7